



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bool

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

1: J Immunol. 1999 Jun 1;162(11):6880-92.

Full Text FREE Links
J Immunol

Human breast carcinoma patients develop clonable oncofetal antigen-specific effector and regulatory T lymphocytes.

Rohrer JW, Barsoum AL, Dyess DL, Tucker JA, Coggin JH Jr.

Department of Microbiology and Immunology, University of South Alabama College of Medicine, Mobile 36688, USA.
jrohrer@jaguar1.usouthal.edu

Oncofetal Ag (OFA) is a 44-kDa glycoprotein expressed during early to mid-gestation fetal development and re-expressed as a surface Ag by tumor cells soon after transformation. The Ag is detectable on all types of human and rodent tumors tested, but is undetectable on normal cells. In experimental animals it is autoimmunogenic and induces potentially protective T cell responses both after experimental immunization and during tumor development subsequent to carcinogenic insult. To determine whether this tumor-associated Ag is also immunogenic for human T lymphocytes, breast carcinoma patients' peripheral blood mononuclear leucocytes were stimulated in vitro with autologous tumor cells in the presence of IL-2, gamma-IFN, and IL-6 for 2 wk. The tumor-reactive cells were then restimulated and cloned by limiting dilution, and the clones were analyzed. We established 24, 19, 11, and 16 tumor-reactive clones from the four respective patients. Of those, 4, 6, 4, and 7, respectively, proliferated specifically to purified OFA. Both CD4 and CD8 OFA-specific clones were established, which responded equally well to purified OFA or 32- to 44-kDa immature laminin receptor protein. All were CD3+, TCR-alpha beta+. All CD4 clones secreted gamma-IFN, but neither secreted IL-4 nor IL-10. Both IFN-gamma-secreting cytotoxic CD8 clones and IL-10-secreting inhibitory CD8 clones were established. Thus, during human cancer development, the same types of OFA-specific effector and regulatory T cells are induced as during murine T lymphomagenesis.

PMID: 10352310 [PubMed - indexed for MEDLINE]

Related Links

Identification of oncofetal antigen/immature laminin receptor protein epitopes that activate BALB/c mouse OFA/iLRP-specific effector and regulatory T cell clones [J Immunol. 2006]

Tumor specific cytolysis by tumor infiltrating lymphocytes in breast cancer. [Cancer. 1994]

CD8 T cell clones inhibit antitumor T cell function by secreting IFN-gamma [J Immunol. 1995]

Characterization of RFM mouse T lymphocyte anti-oncofetal antigen immunity in apparent tumor-free, long-term survivors of sublethal X-irradiation by limiting dilution T lymphocyte cloning. [J Immunol. 1995]

37 kiloDalton oncofetal antigen protein and immature laminin receptor protein are identical, universal T-cell inducing immunogens on primary rodent and human cancers. [Anticancer Res. 1999]
See all Related Articles...

Display AbstractPlus Show 20 Sort By Send to



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search PubMed for

Limits Preview/Index History Clipboard Details

Note: Performing your original search, *oncofetal antigen and epitopes*, in PubMed will retrieve 156 citations.

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

☐ 1: Mol Immunol. 1991 Jul;28(7):703-10.

Links

Epitope analysis of the oncofetal antigen alphafetoprotein using monoclonal antibodies.

Chakraborty M, Mandal C, Mandal C.

Indian Institute of Chemical Biology, Calcutta.

Alphafetoprotein (AFP), an oncofetal antigen, plays very important roles in the early embryonic life and oncogenesis. Under various physiological and pathological conditions AFP exhibits microheterogeneity, probably as a result of differential expression of its epitopes. To analyse the epitopes we have developed a panel of monoclonal antibodies against human AFP purified by a new and efficient method using an immunoadsorbent consisting of polyclonal antibodies immobilized on cyanogen bromide activated Sepharose. Clones producing antibodies of various isotypes, e.g. IgG1, IgG2a, IgG2b, IgA and IgM have been subcloned and characterized. The antibodies showed high avidity for AFP (with half-maximal binding concentrations between 0.012 and 3.87 nM). Mutual inhibition efficiencies of a panel of 14 monoclonal antibodies were determined by RIA. Based on these inhibition data a computer program was used to group these antibodies with respect to their "epitope specificity distance". As a result of this grouping, clones have been identified which can recognize at least five different epitopes on AFP. This panel of antibodies may be very useful for analysis of the epitopic variation of AFP under various physiological and pathological conditions.

PMID: 1713294 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort By Send to

Related Links

Characteristics of five monoclonal anti-alpha-fetoprotein antibodies. [J Biol Chem. 1984]

Monoclonal antibodies against human alpha-fetoprotein. Exploitation of an unusual calcium-dependent interaction with the antigen for analytical and preparative purposes. [J Immunol Methods. 1988]

Monoclonal antibodies to carcinoembryonic antigen: a systematic analysis of antibody specificities by using related normal antigens and evidence for allotypic determinants on carcinoembryonic antigen. [J Immunol. 1984]

Approaches to the production of monoclonal antibodies specific for concanavalin A binding and non-binding forms of alpha-fetoprotein. [J Biol Chem. 1984]

The role of isotype and epitope specificity of monoclonal antibody mixtures in immunodiffusion reactions. [J Immunol. 1984]

See all Related Articles...

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)